

Observations on Chernobyl After 25 Years of Radiophobia

by Zbigniew Jaworowski, M.D., Ph.D., D.Sc.



Vadim Mouchkin/IAEA

Aerial view of the Chernobyl nuclear power plant encased in its sarcophagus.

Ten days after two steam and hydrogen explosions blew up the Chernobyl nuclear reactor, the fire that melted its core died out spontaneously. But the drama of this catastrophe still flourishes, nourished by politics, authorities, media, and interest groups of ecologists, charitable organizations, and scientists. It lives in the collective memory of the world and propagates real health, social, and economic harm to millions of people in Belarus, Russia, and the Ukraine. It is exploited in attempts to strangle the development of atomic energy, the cleanest, safest, and practically inexhaustible means to meet the world's energy needs. The world's uranium resources alone will suffice for the next 470,000 years (IAEA 2008).

Chernobyl was indeed an historic event; it is the only nuclear power station disaster that ever resulted in an occupational death toll, albeit a compar-

tively small one. A vast environmental dispersion of radioactivity occurred that did not cause any scientifically confirmed fatalities in the general population. The worst harm to the population was caused not by radiation, and not to flesh, but to minds.

This catastrophe provided many invaluable lessons. One of them is a recognition of the absurdity of the prevailing linear no-threshold hypothesis (LNT), which assumes that even near-zero radiation dosage can lead to cancer death and hereditary disorders. That the LNT is false, is shown by observing that such damage did not occur after Chernobyl.

Chernobyl was the worst possible catastrophe. It happened in a dangerously constructed nuclear power reactor with a total meltdown of the core and 10 days of free emission of radionuclides into the atmosphere. Probably

The worst possible nuclear plant accident produced no scientifically confirmed fatalities in the general population. But there was enormous political and psychological damage, mainly the result of belief in the lie that any amount of radiation is bad.



Ukrainian Society for Friendship and Cultural Relations with Foreign Countries

A helicopter at the Chernobyl site in 1986, checking the damage to the reactor.

nothing worse could happen. Yet, the resulting human losses, although tragic, were minute in comparison with catastrophes from other energy sources.

Highly sensitive monitoring systems that had been developed in many countries for the detection of fallout from nuclear weapons enabled easy detection of minute amounts of Chernobyl dust, even in remote corners of the world. This added to global epidemics of fear induced by the accident.

Radioactive debris was dispersed into the troposphere and stratosphere of the Northern Hemisphere. up to at least 15 km altitude (Jaworowski and Kownacka 1994). On the first few days after the accident, the concentrations of radiocesium measured at this altitude over Poland (maximum 36.1 mBq/cubic meter at standard temperature and pressure, or STP) was 2 to 6 percent of that at the ground level. Such a high vertical distribution and mixing enabled a small portion of Chernobyl debris to pass over the

equatorial convergence and into the Southern Hemisphere (Philipot 1990), and on to the South Pole (Dibb et al. 1990, Philippot 1990). This was not in agreement with computer-generated models of nuclear accidents, which projected a maximum uplift of fission products to below 3,000 meters altitude (ApSimon et al. 1985, ApSimon and Wilson 1987).

Enormous amounts of radionuclides entered the air from the burning reactor. Yet the total emission was 200 times less than from all of the 543 nuclear warheads exploded in the atmosphere since 1945. The highest estimated radiation dose exposure to the average member of the world population was 0.113 mSv, recorded in 1963 (UNSCEAR 1988). The radiation doses from Chernobyl dust were estimated and compared with natural doses by UNSCEAR (2000a). During the first year after the accident, the average dose received by an average inhabitant of the Northern Hemisphere was estimated by UNSCEAR as 0.045 mSv, that is, less than 2 percent of the average global annual natural dose (2.4 mSv per year).

During the next 70 years, the global population will be exposed to a total Chernobyl dose




Figure 1
WORLDWIDE RADIATION DOSES COMPARED WITH CHERNOBYL

Average annual radiation doses from natural and man-made sources, worldwide, near Chernobyl, and in areas of high natural radiation. The radiation from Chernobyl is dwarfed in comparison with natural background radiation.

Source: Based on UNSCEAR (1988, 1993, 1998, 2000b)

of approximately 0.14 mSv, or 0.08 percent of the natural lifetime dose of 170 mSv. People living in the most contaminated areas of the former Soviet Union received an average individual annual whole-body radiation dose in 1986-1995 of 0.9 mSv in Belarus, 0.76 mSv in Russia, and 1.4 mSv in Ukraine (UNSCEAR 2000b). Average doses estimated for the period 1986-2005 are 2.4 mSv in Belarus, 1.1 mSv in Russia, and 1.2 mSv in Ukraine (UNSCEAR 2008), respectively.

All these doses are dwarfed in comparison with natural radiation doses in some parts of the world. For example, in Ramsar, Iran, natural radiation doses reach more than 400 mSv/year (Mortazawi et al. 2006), and in Brazil and southwestern France, natural radiation doses reach up to more than 700 mSv per year (UNSCEAR 2000b). (See Figure 1.)

A comparison of these doses and epidemiological observations should be a basis of realistic estimates of the latent medical consequences of the Chernobyl accident, rather than risk factors based on the LNT. Such a comparison, and the comparatively minute health consequences, were apparent soon after the catastrophe (Jaworowski 1988), but this information was not shared with the public. Recently the well-known British environmentalist James Lovelock, best known for his Gaia theory, dispelled at length all the usual myths that surround the Chernobyl accident. Lovelock stated that for many years the scientists who could have challenged the nonsense about the catastrophe chose to keep quiet (Murphy 2009). I do not feel guilty.

No harmful health effects have ever been detected in high natural background radiation areas. This is consistent with other studies of the incidence of cancers in exposed populations. In the United States and in China, for example, the incidence of cancers was found to be lower in regions with high natural radiation than in regions with low natural radiation (Frigerio et al. 1973, Frigerio and Stowe 1976, Wei 1990). Among British radiologists exposed mainly to X-rays, cancer mortality was found to be lower by about 50 percent than that in the average male population of England and Wales (Berrington et al. 2001).

Also, in other population groups exposed to low doses of ionizing radiation (i.e., patients diagnosed with iodine-131 and X-rays, dial painters, chemists, and others exposed to ingested or inhaled radium or plutonium, persons exposed to higher levels of indoor radon, and A-bomb survivors) a lower percentage of neoplastic malignancies was observed (Cohen 2000, Luckey 2003, UNSCEAR 1994). A Taiwan study of several thousand residents of apartments contaminated with cobalt-60, who had been chronically exposed to gamma rays for up to 20 years, with total doses estimated to range from 120 to 4,000 mSv, revealed that the cancer mortality and congenital malformations of these residents substantially decreased rather than increased (Chen et al. 2004), suggesting a stimulating or hormetic effect of low doses of low linear-energy-transfer (LET) ionizing radiation.

This finding was partially confirmed by a later study on cancer incidence in a similar Taiwan cohort, in which for all cancers (except leukemia and solid cancers), with the number of cancer cases ranging from 119 to 190, there was a deficit of incidence found in comparison with the unexposed population. In groups of all types of leukemia and of some solid cancers of particular organs, the number of cases was 1 to 2 orders of magnitude smaller than in the first three groups (Hwang 2008).



Environmentalists for Nuclear Energy

An orange from Ramsar, Iran, the region with one of the highest natural background radiation levels in the world. The Ramsar population has lower cancer rates than those of other areas. No harmful health effects have ever been detected in high natural background radiation areas.

About 3,000 reports on radiation hormesis were recently reviewed (Luckey 2003). In one study, among approximately 200,000 American, British, and Canadian nuclear workers exposed to radiation, the total cancer deaths ranged from 27 percent to 72 percent of the total cancer deaths in the control group of non-nuclear workers (Luckey 2003). Such a hormetic deficit invalidates the LNT, because the concept of hormesis transcends any hypothesized dose threshold for excess cancers. If there were no hormesis, the existence of a true threshold for excess cancers might be impossible to demonstrate rigorously, because of the statistical problems of proving an absolute equality of effect in an epidemiological study at a very low dose level. If, however, a deficit of cancers is observed in the population irradiated at a relatively low dose level, as in hormesis, there is often a statistically significant difference at an acceptable confidence level (Webster 1993). This remark of Webster, an UNSCEAR member, reflects discussions in the Committee during preparation of its report on hormesis (UNSCEAR 1994).

A more recent study, based on collective doses for about 400,000 nuclear workers, found a 31 percent decrease in relative cancer mortality (Cardis et al. 2007), but nevertheless concluded that these cancer death data were consistent with the LNT relationship. This conclusion was based on an *ad hoc* accepted assumption of a confounding "healthy worker" effect for the studied cohort. It was assumed that the nuclear workers were selected for employment because they were more healthy. However, the existence of this effect was not supported by their data or by any other factual evidence.

The "healthy worker" effect could be correctly assumed only if the cancer marker diagnostics (ACS 2009) and genetic tests were used in pre-employment screening and selection of these workers. But these procedures were not applied in the Cardis et al. cohort, and even now they are not recommended by the In-



An imaginary six-foot chicken from Chernobyl, written up as news in the National Enquirer in 1986. Other products of hysteria about Chernobyl radiation, including doctored photos, are still in circulation.

ternational Commission on Radiological Protection, the directives of the European Union, or the IAEA International Basic Safety Standards. Thus, this assumption is invalid and explains nothing.

On the other hand, the statistical re-analysis of Cardis et al. data clearly documents that their assumption of a “healthy worker” effect was incorrect, and their data indicated that low doses of ionizing radiation induced a hormetic effect in the exposed nuclear workers (Fornalski and Dobrzynski 2009).

Chernobyl vs. Other Industrial Accidents

In terms of human losses (there were 31 early deaths) the accident in the Chernobyl nuclear power plant was a minor event compared with many other major industrial catastrophes. In the 20th Century, more than 10 such catastrophes have occurred, with tens of thousands of fatalities in each. For example, coal smog killed approximately 12,000 people in London, between December 1952 and February 1953 (Bell and Davis 2001). The annual death toll from accidents in Chinese coal mines reached 70,000 deaths in the 1950s, and 10,000 in the 1990s (WNA 2009). In 1984, about 20,000 people perished after an explosion in a pesticide factory in Bhopal, India (Dhara and Dhara 2002); and the collapse of a hydroelectric dam on the Banqiao river in China in 1975 caused 230,000 fatalities (Altius 2008, McCully 1998, Yi 1998).

The world does not celebrate the anniversaries of these enormous man-made disasters, but year after year we do so for the hundreds and thousands of times less deadly Chernobyl acci-

dent. Ten years ago I discussed the possible causes of this paranoiac phenomenon (Jaworowski 1999). Measured as early deaths per electricity units produced by the Chernobyl facility (nine years of operation, total electricity production of 36 gigawatts of electricity (GWe), 31 early deaths) yields 0.86 deaths/GWe-year). This rate is lower than the average fatalities from a majority of other energy sources.

For example, the Chernobyl rate is 9 times lower than the death rate from liquefied gas (Hirschberg et al. 1998) and 47 times lower than from hydroelectric stations (40.19 deaths/GWe-year including the Banqiao disaster). But the political, economic, social, and psychological impact of Chernobyl was enormous. Let’s examine what happened starting with my personal experience.

Psychology Tuned by LNT

At about 9 A.M. on Monday, April 28, 1986, at the entrance to my institute in Warsaw, I was greeted by a colleague who said: “Look, at 7:00 we received a telex from a monitoring station in northern Poland saying that the beta radioactivity of the air there is 550,000 times higher than the day before. I found a similar increase in the air filter from the station in our backyard, and the pavement here is highly radioactive.”

This was a terrible shock. My first thought was, A NUCLEAR WAR! It is curious that all my attention was concentrated on this enormous rise of total beta activity in the air used to monitor radiation emergencies from nuclear test fallout. Many years spent during the Cold War on preparations to defend the Polish population against the effects of a nuclear attack had conditioned my colleagues and me to have such an exaggerated reaction.

We reacted that way although we knew, that on this first day of Chernobyl in Poland, the dose rate of external gamma radiation penetrating our bodies was higher only by a factor of 3 from the day before, and it was similar to the average natural radiation doses which from time immemorial we have received from ground and cosmic radiation. At 11 A.M., after we had collected enough dust from the air for gamma spectrometry measurements, we discovered that it contained cesium-134. Thus, we knew that its source was not an atomic bomb, but a nuclear reactor. This was tranquilizing news, which did not, however, calm our frantic behavior.

In 1986, the impact of a dramatic increase in atmospheric radioactivity dominated my thinking—and everybody else’s. This state of mind led to immediate consequences. First there were various hectic actions, such as *ad hoc* coining of different limits for radionuclides in food, water, and other things. In particular countries, these limits varied by a factor of many thousands, reflecting various political and mercenary factors and the emotional states of the decision makers.

For example, Sweden allowed for 30 times more radioactivity in imported vegetables than in domestic ones, and Israel allowed less radioactivity in food from Eastern Europe than from Western Europe. The cesium-137 concentration limit in vegetables imposed in the Philippines was 22 Bq per kg, 8,600 times lower than in the more pragmatic United Kingdom (Salo and Daglish 1988). In Poland, a group of nuclear physicists and engineers proposed a cesium-137 limit of 27 Bq in 1 kilogram for any kind of food, but, fortunately, the authorities decided more soberly and imposed a 1,000 Bq limit.

Behind these restrictions, meaningless from the point of view of human health, stood three factors: (1) emotion; (2) the LNT mindset and the international recommendations based on it; and (3) a social need to follow an old medical rule, *Ut aliquit fecisse videatur* (to make it appear that something is being done). That third factor was a placebo used by the authorities to dodge the worst kind of criticism, i.e., accusations of inactivity in the face of a monstrous disaster. This led to an overreaction in Europe and in some other countries, but at the greatest scale and with the most severe consequences in the Soviet Union.

The High-Cost of Hysteria

The costs of these regulations were enormous. For example, Norwegian authorities introduced a cesium-137 concentration limit of 6,000 Bq/kg in reindeer meat and game, and a 600 Bq/kg limit for sheep (Henriksen and Saxebol 1988). A Norwegian eats an average of 0.6 kg of reindeer meat per year. The average radiation dose from eating this amount of meat is estimated to be about 0.047 mSv per year. Thus, this measure was aimed to protect Norwegians against a radiation dose about 200 times lower than the natural dose in some regions of Norway of 11 mSv per year (UNSCEAR 1982).

The costs of this protection climbed to over \$70 million in 1986, and in the 1990s it was still about \$4 million per year (Christensen 1989, Idas and Myhre 1994). This means that unnecessary and wasteful restrictions, once implemented under the influence of the above three factors, have a long lifetime.

The hysterical reaction of authorities, further excited by extremely exaggerated media reports, is well exemplified by the Japanese government's cancellation of a several-hundred-million (in U.S. dollars) contract for shipping Polish barley to Japan for the production of beer. This happened in May 1986, a few days after completely false information of extreme contamination of Poland by Chernobyl fallout appeared on the front page of the biggest Japanese daily, *Asahi Shimbun*. It screamed with block letters, "DUST OF DEATH IN POLAND," and it cited my name as the source of the information.

I was asked by the Polish government to write a text in English which might be used to avert this loss of money. I did this during a weekend spent with my wife in our cottage on the banks of the Vistula, together with John Davis, the American ambassador to Poland, and his charming wife, Helene. When I finished my writing assignment, I asked John to correct the language. He said that the English was almost OK, but not exactly in proper diplomatic style. He then proceeded to change the text completely.



Katarzyna Dopieralska-Skowronska

Author Zbigniew Jaworowski, speaking here at a 2005 geophysical meeting in Warsaw.

On Monday a spokesman for the communist government asked me to read the text at his press conference. I presented the talk, but after I finished, he distributed copies of the talk to the waiting flock of journalists. He was totally unaware that the written text had been prepared by the U.S. ambassador. A visit by the Japanese ambassador to our Central Laboratory for Radiation Protection managed to salvage the contract.

A few days later, Ambassador Davis arranged an international deal for shipment by air of large quantities of powdered milk for Polish children, to replenish strategic reserves that were rapidly being depleted. This was not an easy task, because other European countries, in a similar position to ours, refused to sell their milk. As we now know, during the next four years the Davises played a delicate but pivotal role in realizing a major goal for the people of Poland, the Solidarity movement's victory over communism (Davis 2009, Davis et al. 2006). As explained below, Solidarity's triumph was related to the Chernobyl accident.

The Costly Folly of LNT

A classic example of wastefully applying the LNT principle to the Chernobyl emergency was provided by Swedish radiation-protection authorities. When the farmers near Stockholm discovered that the Chernobyl accident had contaminated their cows' milk with cesium-137, above the limit of 300 Bq per liter imposed by authorities, they wrote the authorities to ask if their milk could be diluted with uncontaminated milk from other regions, to bring it below the limit. This would be done by mixing 1 liter of contaminated milk with 10 liters of clean milk.

To the farmers' surprise and disappointment, the answer was "no," and the milk was then to be discarded. This was a strange ruling since it has always been possible to reduce pollutants to safer levels by dilution. We do this for other pollutants in food-stuffs, and we dilute fumes from fireplaces or ovens with atmospheric air in the same way that nature dilutes volcanic emissions or forest fire fumes. The Swedish authorities explained that even though the individual risk could be reduced by diluting the milk, this would, at the same time, increase the number of consumers. Thus, the risk would remain the same, but now spread over a larger population (Walinder 1995).

Although ridiculous, this was a faithful application of the International Commission on Radiological Protection recommendations, based on the LNT assumption and its offspring, the concept of "collective dose"; that is, reaching terrifyingly large numbers of man-sieverts by multiplying tiny, innocuous individual radiation doses by a large number of exposed people.



Elisabeth Zeiller/IAEA

A local market, where food samples were taken for use in the IAEA diet study of the Chernobyl Assessment Project. The hysteria around the accident and the adherence to the LNT thesis led to widely varying regulations restricting food use that cost European nations millions of dollars.

In an earlier paper, I exposed the negative consequences and lack of sense in the LNT assumption, and the collective dose and dose-commitment concepts (Jaworowski 1999). The application of these principles has caused the costs of the Chernobyl accident to exceed \$100 billion in Western Europe (Becker 1996), and much more in post-Soviet countries where it has led to untold suffering and the pauperization of millions of people. The international institutions standing behind this assumption and these concepts certainly will not admit responsibility for their disastrous consequences. They should.

Some LNT History

The linear no-threshold hypothesis was accepted in 1959 by the International Commission on Radiological Protection (ICRP 1959) as the philosophical basis for radiological protection. This decision was based on the first report of the newly established United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR 1958). A large part of this report was dedicated to a discussion of linearity and of the threshold dose for adverse radiation effects.

Fifty years ago, UNSCEAR's stand on this subject was formed after an in-depth debate that was not without influence from the political atmosphere and issues of the time. The Soviet, Czechoslovakian, and Egyptian delegations to UNSCEAR strongly supported the LNT assumption, and used it as a basis for recommendation of an immediate cessation of nuclear test explosions. The LNT was also supported by the Soviet Union during the later years of the Cold War (Jaworowski 2009), and this was consistent with the thinking of American authorities.

The target theory prevailing in the 1950s and the then new results of genetic experiments with fruit flies irradiated with high doses and dose rates, strongly influenced this debate. In 1958, UNSCEAR stated that contamination of the environment by nuclear explosions increased radiation levels all over the world and thus posed new and unknown hazards for present and future generations. These hazards, UNSCEAR stated, cannot be controlled, and "even the smallest amounts of radiation are liable to cause deleterious genetic, and perhaps also somatic, effects."

This sentence had an enormous impact in subsequent decades, and has been repeated in a plethora of publications. Even today, it is taken as an article of faith by the public. However, throughout the entire 1958 report, the original UNSCEAR view on LNT remained ambivalent. As an example, UNSCEAR accepted as a threshold for leukemia a dose of 4,000 mSv (page 42); but at the same time, the Committee accepted a risk factor for leukemia of 0.52 percent per 1,000 mSv, assuming LNT (page 115). The committee quite openly presented this difficulty, and showed its consequences in a table (page 42).

Continuation of nuclear weapons tests in the atmosphere was estimated to cause 60,000 leukemia cases worldwide, if no threshold were assumed, and zero leukemia cases if a threshold of 4,000 mSv were in place. In its final conclusions, UNSCEAR pinpointed this dilemma: "Linearity has been assumed primarily for purposes of simplicity," and "There may or may not be a threshold dose. The two possibilities of threshold and no-threshold have been retained because of the very great differences they engender."

After a half-century, we still discuss the same problem. In 1958, UNSCEAR had no doubts about major genetic defects in the world population that could be caused by nuclear test fallout, and estimated them as high as 40,000. But later, the Committee learned that even among the children of highly irradiated survivors of atomic bombings, no statistically significant genetic damage could be demonstrated (UNSCEAR 2001).

However, in the International Commission on Radiological Protection document of 1959, no such controversy and no hesitations appeared. The LNT was arbitrarily assumed, and serious epistemological problems related to the impossibility of finding harmful effects at very low levels of radiation were ignored. Over the years, the working assumption of the International Commission in 1959 came to be regarded as a scientifically documented fact by the mass media, public opinion, and even many scientists. The LNT assumption, however, is not a proven scientific principle, and belongs in the realm of administration (Jaworowski 2000).

LNT *ad Absurdum*

The absurdity of the LNT was brought to light in 1987, when minute doses of Chernobyl radiation were used to calculate

that 53,000 people would die of Chernobyl-induced cancers over the next 50 years (Goldman et al. 1987). This frightening death toll calculation was derived simply by multiplying the trifling Chernobyl doses in the United States (0.0046 mSv per person) by the vast number of people living in the Northern Hemisphere, and by a cancer risk factor based on epidemiological studies of 75,000 atomic bomb survivors in Japan.

But the A-bomb survivor data are irrelevant to such estimates because of the difference in the individual doses and dose rates. A-bomb survivors were flashed within less than a second by radiation doses at least 50,000 times higher than any dose that U.S. inhabitants will ever receive over a period of 50 years from the Chernobyl fallout.

We have reliable epidemiological data for a dose rate of perhaps 1,000 or 6,000 mSv per second in Japanese A-bomb survivors. But there are no such data for human exposure at a dose rate of 0.0045 mSv over 50 years, nor will there ever be any. The dose rate in Japan was larger by a factor of about 10^{12} than the Chernobyl dose rate in the United States. Extrapolating over such a vast span is neither scientifically justified nor epistemologically acceptable. It is also morally suspect (Walinder 1995). Indeed, Lauriston Taylor, the late president of the U.S. National Council on Radiological Protection and Measurements, deemed such extrapolations to be a “deeply immoral use of our scientific heritage” (Taylor 1980).

In its document on protection of the public in a major radiation emergency, the International Commission on Radiological Protection recommended the administration of stable iodine, in the form of tablets to be taken before, or as soon as possible after, the start of exposure to radioactive iodine-131 (ICRP 1984). The Commission advised applying this prophylactic measure to everybody—pregnant women, neonates, young infants, and adults—starting at the projected thyroid dose of 50 mSv. This recommendation was based on the LNT dogma. We followed it in Poland.

In the late afternoon of April 28, 1986, we learned from the BBC that there was a reactor accident in Chernobyl. We had seen the radioactive cloud flowing over Poland from east to west, and we had the first data on concentration levels of radioiodine in grass and soil in eastern Poland and in Warsaw. Using these data, I calculated that contamination of thyroid glands of Polish children might reach a limit of 50 mSv, and much more if the situation in Chernobyl and weather conditions further aggravated the situation.

Meaningless Administration of Stable Iodine

In our Institute we had no information from the Soviet Union on the current state of affairs or of any projections regarding the behavior of the destroyed reactor. Therefore, we assumed that in the next few days the radioactivity in the air would increase and cover the whole country. We prepared a portfolio of coun-



Courtesy of Zbigniew Jaworowski

Prophylactic doses of stable iodine were administered in liquid form (Lugol) within three days of the Chernobyl accident to 18.5 million children and adults in Poland. In agreement with the recommendations of the International Commission on Radiological Protection and the International Atomic Energy Agency (all based on the LNT), the author had recommended this vast operation to the Polish government. Now, he regards this action as futile.

termeasures to be implemented by the government.

I presented this project at a meeting of the deputy prime minister, several ministers, and high ranking secretaries of the Central Committee of the Polish United Workers Party, at about 4 A.M. on April 29. The most important measure recommended—and also accepted after a short discussion by this mixture of government and party officials—was stable iodine prophylaxis to protect the thyroid glands of children against iodine-131 irradiation.

Administration of stable iodine in liquid form (as a solution of Lugol) was initiated in the northeastern part of Poland, approximately 38 hours after we discovered the Chernobyl fallout (at approximately midnight on April 28). Treatment was given for the next three days, and about 18.5 million people, including adults, received the stable iodine drug.

We were able to perform this action successfully because we had already made plans for implementing nuclear war emergency measures. In the 1960s, our Institute had recommended that the government prepare for such an event by distributing strategic stores of stable iodine at sites all over the country, as the only reasonable measure against body contamination from fission products. The program was implemented in the early 1970s, and each Polish pharmacy, hospital, and various other institutions had large supplies of iodine.

At the time of the Chernobyl accident, Poland had more than enough iodine ready for use for approximately 100 doses for each Polish citizen. A few years after the catastrophe, it was estimated that in the more contaminated parts of the country the average thyroid radiation dose in the 1- to 10-year-old age group was about 70 mSv, and in about 5 percent of children the maximum dose was about 200 mSv (Krajewski 1991).



Petr Pavlicek/IAEA

The ghost town of Prip'yat in July 2005. Its 47,000 residents, including 17,000 children, were completely evacuated the day after the accident in 1986. Prip'yat was built in the 1970s to house Chernobyl workers in the 1970s, it was one of the “youngest” towns in the then Soviet Union; the average age of its inhabitants was 26. Today, it is frozen in time. The ferris wheel (center left) is part of an amusement park that never opened. It should be resettled!

A decade later, we learned that among the more than 34,000 Swedish patients who were not suspected of having thyroid cancers, and whose thyroids were irradiated with iodine-131 up to doses of 40,000 mSv (average dose 1,100 mSv), there was no statistically significant increase in thyroid cancers, but rather a 38 percent decrease in their incidence (Dickman et al. 2003, Hall et al. 1996, Holm et al. 1988).

If I knew then what I know today, I would not have recommended to the Polish government such a vast prophylactic action, not because of its allegedly adverse medical effects—there were none (Nauman 1989)—but because its practical positive health effect was meaningless.

Harmful Mass Evacuations

The most nonsensical, expensive, and harmful action, however, was the evacuation of 336,000 people from contaminated regions of the former Soviet Union, where the radiation dose from Chernobyl fallout was about twice the natural dose. Later, this limit was decreased to even below the natural level, and was some five times lower than the radiation dose rate of 5.25 mSv/year at Grand Central Station in New York City, which is constructed with natural granite (Benenson et al. 2006).

Contaminated areas were defined as being those where the average cesium-137 ground deposition density exceeded 37 kBq per square meter. In the Soviet Union, these areas covered 146,100 square kilometers. The Chernobyl fallout of about 185 kBq per square meter or more also covered large areas of Austria, Bulgaria, Finland, Norway, and Sweden (UNSCEAR 2000b). Small areas with Chernobyl fallout, reaching up to about 185 kBq per square meter, were also found in other countries (Great Britain, Greece, Romania, Switzerland, and Turkey (EUR 1996)).

The average radiation doses received in areas with a cesium-137 deposition density of about 37 kBq per square meter were estimated at about 1.6 mSv during the first year after the Chernobyl accident, and the lifetime dose (after 70 years) was predicted to reach 6 mSv (UNSCEAR 1988). This activity level is 10 times lower than the average amount (400 kBq per square meter) of about 50 natural radionuclides present in a 10-cm-thick layer of soil (Jaworowski 2002). The corresponding Chernobyl lifetime radiation dose is 28 times lower than the average natural lifetime dose of about 170 mSv. But the annual dose from 37 kBq of cesium-137 per square meter was similar to the 1 mSv/year dose limit recommended by the International Commission on Radiological Protection for the general population, and this is why it was accepted by the Soviet authorities as a yardstick for remedial measures.

The evacuation caused great harm to the populations of Belarus, Russia, and the Ukraine. It led to mass psychosomatic disturbances, great economic loss and traumatic social consequences. According to Academician Leonid A. Ilyin, the leading Russian authority on radiation protection, the mass relocation was implemented by the Soviet government under the pressure of populists, ecologists, and self-appointed specialists, and it was done against the advice of the best Soviet scientists (Ilyin 1995, Ilyin 1996). The really dangerous air radiation dose rate of 1 Gy/hour on April 26, 1986 (0.01 Gy/hour two days later) covered an uninhabited area of only about 0.5 square kilometers in two patches, reaching up to a distance of 1.8 km southwest of the Chernobyl reactor (UNSCEAR 2000b).

Based on these data, there was no valid reason for the mass evacuation of 49,614 residents from the city of Prip'yat and the village of Yanov, situated about 3 km from the burning reactor. In these settlements, the radiation dose rate in the air on April 26, 1986 was 1 mSv/hour (UNSCEAR 2000b), and two days later it was only 0.01 mSv/hour. Thus, with a steadily decreasing radioactivity fallout, the dose rate was not dangerous at all.

However, according to L.A. Ilyin, one of the leaders of the Chernobyl rescue team, there was a danger that the corium (the melted core of the reactor, with a total volume of about 200 cubic meters, a mass of about 540 tons, and a temperature of about 2000°C,) might penetrate down through the concrete floor and spread to rooms below. The team suspected that in these rooms there could have been a great volume of water, with which the corium could come into contact. This would have led to a much more powerful explosion than the initial one, and caused a vastly greater emission of radioactivity, which could have covered Prip'yat and Yanow with lethal fall-



Fornalski 2009

Radiation measurement in Pripyat on April 10, 2008 at a sports stadium in the downtown area of the abandoned city, which is about 4 km northwest from the Chernobyl reactor. The dose rate was 0.28 $\mu\text{Sv}/\text{hour}$ or 2.5 mSv/year . This is more than 10 times lower than the natural radiation in many areas of the world.

out. Therefore, the evacuation of the whole population of these localities was a correct precautionary measure that was carried out in an orderly manner in only two hours.

But the evacuation and relocation of the remaining approximately 286,000 people, of whom there were about 220,000 after 1986 (UNSCEAR 2000b), was an irrational overreaction, induced in part by the influence of the International Commission of Radiological Protection and International Atomic Energy Agency recommendations based on the LNT (Ilyin 1995). The current reluctance of the Ukrainian authorities to resettle the residents back to Pripyat (now a slowly decaying ghost town and tourist attraction) does not seem rational. The radiation dose rate measured on April 10, 2008 in the streets of this city ranged from 2.5 to 8.4 mSv/year , i.e., more than 10 times lower than natural radiation in many regions of the world (Fornalski 2009).

Psychosomatic Epidemics

In addition to the 28 fatalities among rescue workers and employees of the power station, caused by very high doses of radiation (2.9-16 Gy), and 3 deaths due to other reasons (UNSCEAR 2000b), the only real adverse health consequences of the Chernobyl catastrophe among approximately 5 million people living in the contaminated regions were the epidemics of psychosomatic afflictions that appeared as diseases of the digestive and circulatory systems and other post-traumatic stress disorders, such as sleep disturbance, headache, depression, anxiety, escapism, learned helplessness, unwillingness to cooperate, over-dependence, alcohol and drug abuse, and suicides.

These diseases and disturbances could not have been caused by the minute irradiation doses from the Chernobyl fallout (average dose rate of about 1 to 2 mSv/year), but they were caused by radiophobia, a deliberately induced fear of radiation, aggravated by wrongheaded administrative decisions and even, paradoxically, by increased medical attention, which leads to diagnosis of subclinical changes that persistently hold the

attention of the patient.

Bad administrative decisions made several million people believe that they were *victims of Chernobyl*, although the average annual dose they received from Chernobyl radiation was only about one third of the average natural dose. This was the main factor responsible for the economic losses caused by the Chernobyl catastrophe, estimated to have reached \$148 billion by 2000 for the Ukraine, and to reach \$235 billion by 2016 for Belarus.

Psychological factors and a failure to teach radiological protection in medical school curricula might have led to abortions of wanted pregnancies in Western Europe during the period soon after the accident, where physicians wrongly advised patients that Chernobyl radiation posed a health risk to unborn children. However, numerical estimates of this effect (Ketchum 1987, Spinelli and Osborne 1991) cast doubt on this assumption.

Similarly uncertain are estimates of the number of decisions against conception probably taken in Europe during the first few months after the accident (Trichopoulos et al. 1987). This problem was discussed in 1987 by an IAEA Advisory Group, which concluded that medical practitioners having direct contact with the population at large are among the most important persons who might develop the right perception of risks in nuclear emergencies, prevent social panic and overreactions, and help to ensure the rational behavior in the society.



Petr Pavlicek/IAEA

A doctor from the IAEA International Chernobyl Project examines a child in Ukraine, 1990. Although the average radiation dose to the several million people around Chernobyl was only about one third of the average annual dose from natural radiation, the panic and radiophobia after the accident created a class of "Chernobyl victims," with many disorders related to radiophobia, not actual radiation dose.



Elisabeth Zeiler/IAEA

A group of villagers being interviewed for the IAEA epidemiological study during the International Chernobyl Assessment Project.

After the Chernobyl accident the public very often turned for help to medical practitioners, but physicians were unable to provide realistic advice, even on minor problems. This was because medical curricula did not at that time prepare doctors for nuclear emergencies. In none of the nine countries represented at the meeting were the principles of radiobiology and radiation protection included in medical school curricula (IAEA 1987). Lack of knowledge in this important group was among the factors that increased public anxiety and stress. It seems that now, two decades later, the situation in this respect is very much the same.

Effects of Chernobyl Fallout on the Population

In 2000, the United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR 2000b) and in 2006, the United Nations Chernobyl Forum (a group composed of representatives from eight U.N. organizations, the World Bank, and the governments of Belarus, Russia, and the Ukraine) stated in their documents that, except for thyroid cancers in the population of highly contaminated areas, there was no observed increase in the incidence of solid tumors and leukemia, and no observed increase in genetic diseases. An increase in registration of thyroid cancers in children under 15 years old was first found in 1987, one year after the accident, in the Bryansk region of Russia, and the greatest incidence, of 0.027 percent of children under 15 was found in 1994.

Both of these studies were made too early to be in agreement with what we know about radiation-induced cancers. The mean latency period for ma-

lignant thyroid tumors in adults and children exposed to external and internal medical irradiation with less than 20 to more than 40 Gy is about 28 years (Kikuchi et al. 2004, UNSCEAR 2000b).

Kikuchi et al. tried to explain the discrepancy between the clinical experience and the Chernobyl findings with some exotic ideas, such as, for example, radiation leakage or other environmental conditions; exposure to carcinogens that occurred near Chernobyl prior to the nuclear accident; and a genetic predisposition of the population to thyroid cancer. However, the serendipitous effect of mass screening and diagnosis, already suspected in 1987, is a more likely explanation.

The Clinical Screening Effect

The number of 4,000 new thyroid cancers registered among the children from Belarus, Russia, and the Ukraine should be viewed in the context of the extremely high occurrence of these dormant subclinical malignant tumors that contain transformed tumor cells, which are quite common in the world population (Akslen and Naumov 2008, Weinberg 2008). For example, the incidence of occult thyroid cancers, varies from 5.6 percent in Colombia, 9.0 percent in Poland, 9.3 percent in Minsk (Belarus), 13 percent in the United States, and 28 percent in Japan, to 35.6 percent in Finland (Harach et al. 1985, Moosa and Mazzaferri 1997). In Finland, these dormant thyroid cancers are observed in 2.4 percent of children (Harach et al. 1985), that is, some 90 times more than the maximum observed in the Bryansk region, the most contaminated in Russia.

In Minsk, Belarus, the normal incidence of occult thyroid cancers is 9.3 percent (Furmanchuk et al. 1993). The "Chernobyl" thyroid cancers are of the same histological type and are similar in invasiveness to the occult cancers (Moosa and Mazzaferri 1997, Tan and Gharib 1997). Since 1995, the number of regis-



Petr Pavlicek/IAEA

The new town of Slavutich, 50 km from Chernobyl, which was built for the displaced persons of Chernobyl.

tered cancers has tended to decline. This is not in agreement with what we know about radiation-induced thyroid cancers, whose latency period is about 5-10 years after irradiation exposure (Inskip 2001), and whose risk increases until 15-29 years after exposure (UNSCEAR 2000a).

In the United States the incidence rate of thyroid tumors detected between 1974 and 1979 during a screening program, was 21 times higher than before the screening (Ron et al. 1992), an increase similar to that observed in three former Soviet countries. It appears that the increased registration of thyroid cancers in contaminated parts of the countries affected by Chernobyl is a classical screening effect.

According to the regulations of the Belarusian Ministry of Health, the thyroids of all people who were younger than 18 in 1986 and those of each inhabitant of contaminated areas must be diagnosed every year (Parshkov et al. 2004). More than 90 percent of children in contaminated areas are now examined for thyroid cancers every year with ultrasonography and other methods. It is obvious that such a vast-scale screening, probably the greatest in the history of medicine, resulted in finding thousands of the occult cancers, or incidentalomas, expanded to forms detectable by modern diagnostic methods that were not in routine use in the Soviet Union before 1986.

Data for the past 20 years, published by Ivanov et al. in 2004 and cited in the UNSCEAR and Chernobyl Forum documents (Forum 2005, Forum 2006, Ivanov et al. 2004, UNSCEAR 2008) show, in comparison to the Russian general population, that there was a 15 to 30 percent lower mortality from solid tumors among the Russian Chernobyl emergency workers, and a 5 percent lower average solid tumor incidence among the population of the Bryansk district, the most contaminated in Russia (Figures 2 and 3).

In the most exposed group of these people (with an estimated average mean radiation dose of 40 mSv), a 17 percent decrease in the incidence of solid tumors of all kinds was found. In the Bryansk district, the leukemia incidence is not higher than in the Russian general population. According to UNSCEAR (2000b), no increase in birth defects, congenital malformations, stillbirths, or premature births could be linked to radiation exposures caused by the Chernobyl fallout. The final conclusion of the UNSCEAR 2000 report is that the population of the three main contaminated areas, with a cesium-137 de-




Figure 2
STANDARD MORTALITY RATIOS FOR SOLID CANCERS AMONG THE RUSSIAN EMERGENCY WORKERS

The values of standard mortality ratios (SMR) indicate how the cancer mortality of emergency workers differs from that of the general population of Russia, which was used as a control group (SMR = 1.0). The deficit of cancers among these workers between 1990 and 1999, ranged between 15 percent and 30 percent.

Source: Ivanov et al. 2004, p. 225




Figure 3
STANDARD INCIDENCE RATIOS FOR SOLID CANCERS IN THE BRYANSK REGION OF RUSSIA

The average deficit of cancers in the inhabitants of the Bryansk region was 5 percent, and in the most exposed group (mean radiation dose of 40 mGy) it was 17 percent.

Source: Ivanov et al. 2004, pp. 373-374



IAEA

Checking radiation in a house in the village of Babovichi, Russian Federation, in August 1990.

position density greater than 37 kBq/square meter, need not live in fear of serious health consequences, and forecasts that generally positive prospects for the future health of most individuals should prevail.

The publications of the U.N. Chernobyl Forum present a rather balanced overview of the Chernobyl health problems, but with three important exceptions. The first (mainly after Cardis et al. 2005) is ignoring or downplaying the effect of screening for thyroid cancers in about 90 percent of the population (see discussion above), and interpreting the results with a linear no-threshold dose-response model.

The paper by Cardis et al., however, was criticized for this interpretation, as not confirmed by the data presented and attributing most of the thyroid cancers to radiation (Scott 2006). Both the Chernobyl Forum and the 2005 and 2006 papers by Cardis et al. ignore the aforementioned fundamental problem of occult thyroid cancers in the former Soviet Union and elsewhere in Europe.

The incidence of thyroid occult cancers increased rapidly after the advent of new ultrasonography diagnostics (Topliss 2004), reaching up to 35.6 percent (see above). This incidence is more than 1,300 times higher than the maximum thyroid cancer incidence found in the Bryansk region of Russia in 1994 (UNSCEAR 2000b), which implies a vast potential for bias. It seems that there still has not been an epidemiological study of the temporal changes of intensity of thyroid screening in the former Soviet Union. The conclusions of the epidemiological studies that did not take into account these changes in screening may be invalid.

In the Bryansk region of Russia, the thyroid cancer incidence




Figure 4

CHERNOBYL RADIATION AFTER THE ACCIDENT

The radiation dose rate in air on April 26, 1986 in the local area of the Chernobyl reactor. Units of the isolines are sieverts per hour. Only in the two patches inside the 1 Sv isolines were the dose rates life endangering, during the first two days. After two days, the dose rates decreased about 100 times.

Source: Adapted from UNSCEAR 2000




Figure 5

RADIATION PLUMES AFTER THE CHERNOBYL ACCIDENT

The plumes of radiation formed by meteorological conditions on the days following March 26, 1986. The dates and times are indicated in Greenwich Mean Time.

Source: Adapted from UNSCEAR 2000

was found to be 45 percent higher in males and 90 percent higher in females, than for the Russian population as a whole. However, when dose-response analyses were performed, using external and internal comparisons, no positive association of thyroid cancers with radiation dose was observed. Instead, a negative association was observed, i.e. a hormetic effect (Iva-




Figure 6
SURFACE GROUND MAP OF CESIUM-137
RELEASED IN THE CHERNOBYL ACCIDENT

Source: Adapted from UNSCEAR 2000

nov et al. 2004). These results strongly suggest that the increased cancer rates in Bryansk (and, by implication, in other contaminated regions) compared with general population rates are the result of thyroid cancer screening and better reporting, rather than radiation exposure (Ron 2007).

Even more important a problem in the U.N. Chernobyl Forum report was that it ignored the decrease of thyroid cancer incidence of up to 38 percent, after the iodine-131 treatment of many thousands of non-cancer patients with thyroid radiation doses similar to, or higher than, those from the Chernobyl fallout (Dickman et al. 2003, Hall et al. 1996, Holm et al. 1991, and Holm et al. 1988).

The second problem with the Chernobyl Forum report is estimation of deaths among the patients with acute radiation disease. From among 134 persons with this disease who had been exposed to extremely high radiation doses, 31 died soon after the accident. Among the 103 survivors, 19 died before 2004. Most of these deaths were caused by such disorders as lung gangrene, coronary heart disease, tuberculosis, liver cirrhosis, fat embolism, and other conditions that can hardly be defined as caused by ionizing radiation. Nevertheless, the Chernobyl Forum presents them as a resulting from high irradiation and sums them up to arrive at a total of approximately 50 victims of acute irradiation.

After many summers, all the 103 survivors will eventually die. The Chernobyl Forum philosophy would then count them all, yielding a round total of 134 victims of high irradiation. In fact, the mortality rate among these 103 survivors was 1.08 percent per year, that is, less than the average mortality rate of 1.5 percent in the three affected countries in 2000 (GUS 1991).

And finally, the third Chernobyl Forum “problem” is its projections of future fatalities caused by low-level Chernobyl radiation, from 4,000 up to exactly 9,935 deaths. These numbers are

not based on epidemiological data of cancer mortality observed during the past 20 years by Ivanov et al. No such increase was demonstrated by Ivanov et al. (2004), but rather a decrease of solid tumor and leukemia deaths among exposed people. These epidemiological data, rather than the LNT assumption, should be used as the basis for a realistic projection of the future health of the millions of people officially labeled “victims of Chernobyl.”

However, the Chernobyl Forum instead chose to use the LNT radiation risk model (ICRP 1991) and performed a simplistic arithmetical exercise, multiplying small doses by a great number of people, and including a radiation risk factor deduced from the Hiroshima and Nagasaki studies.

This is an entirely fallacious method. People living in areas highly contaminated by the Chernobyl fallout were irradiated during a protracted time. The dose

rates in Hiroshima and Nagasaki, in contrast, were higher by a factor of about 10^{11} than the average dose rate of the Chernobyl victims that was used in the Forum’s projections. The result of this exercise is nothing more than a fantastic lie.

Several scientific and radiation protection bodies, including UNSCEAR, the Health Physics Society (Mossman et al. 1996), the French Academy of Science (Tubiana 1998), and even the chairman of the International Commission on Radiological Protection (Clarke 1999), advised against making such calculations. Merely publishing these numbers is harmful and petrifies the Chernobyl fears.

Any efforts to explain the intricacies of radiation risk assessments to the public, or to compare these numbers with the much higher level of spontaneous cancer deaths, will be futile exercises. The past 20 years has proved that such efforts are worthless. Making such calculations keeps a lot of people busy and well, but has no relationship to reality and honesty. The Forum’s elucubrations, however, pale in comparison with recent estimates by other bodies such as Greenpeace (Greenpeace 2006, Vidal 2006), predicting the incidence of millions of Chernobyl cancers and hundreds of thousands of deaths.

Remove the Chernobyl Restrictions!

It is reassuring, however, that 16 years after the Chernobyl catastrophe, another group, composed of four U.N. organizations—the United Nations Development Programme (UNDP), the World Health Organization (WHO), the U.N. International Children’s Emergency Fund (UNICEF) and the U.N. Office for the Coordination of Humanitarian Affairs (UNOCHA)—dared to state in its 2002 report, based on UNSCEAR studies, that a great part of the billions of dollars used to mitigate the consequences of the Chernobyl accident was spent incorrectly. The

dollars spent in these efforts did not improve, but actually worsened, a deteriorating situation for 7 million so-called “victims of Chernobyl” and solidified the psychological effects of the catastrophe and the wrong decisions of the authorities.

The report (UNDP 2002) recommended that the three post-Soviet countries and the international organizations abandon the current policy. The misguided basis of this policy, i.e. expectation of mass radiation health effects, was responsible for the enormous and uselessly expended resources sacrificed for remediation efforts. Instead, the report presented 35 practical recommendations needed to stop the vicious cycle of Chernobyl frustrations, social degradation, pauperization, and the epidemic of psychosomatic disorders. The recommendations suggest a reversal of the policy of concentrating attention on nonexistent radiation hazards, and propose that relocated individuals be allowed to return to their old settlements. That is, that essentially all of the restrictions should be removed.*

But here we enter a political mine-field. How well will people accept losing the mass benefits (equivalent to about \$40 a month) that they poetically call a “coffin bonus”? How can it be explained to them that they were made to believe that they were the “victims” of a nonexistent hazard; that the mass evacuations were an irresponsible error; that for 20 years, people were unnecessarily exposed to suffering and need; that vast areas of land were unnecessarily barred from use; and that their countries’ resources were incredibly squandered?

One can read in many publications that the Chernobyl catastrophe had serious political implications and was an important factor in the dismantling of the Soviet Union and in attempts to control nuclear arms. As Mikhail Gorbachev stated:

The nuclear meltdown at Chernobyl 20 years ago even more than my launch of perestroika, was perhaps the real cause of the collapse of the Soviet Union five years later. Chernobyl opened my eyes like nothing else: it showed the horrible consequences of nuclear power. One could



Petr Pavlicek/IAEA

A farmer in Jelno, July 2005. Jelno is a village 300 km from Chernobyl, which was affected by contamination from the accident because of weather conditions. Now the population has gone back to the land. “Social upheaval, however,” the IAEA noted, “has left farmers with only primitive tools of the trade.”



Petr Pavlicek/IAEA

“Jelno is a town where time has stood still,” the IAEA noted, unlike the new settlement of Slavutich.

now imagine much more clearly what might happen if a nuclear bomb exploded—one SS-18 rocket could contain a hundred Chernobyls. Unfortunately, the problem of nuclear arms is still very serious today (Gorbachev 2006).

Would fulfilling the recommendations of the United Nations Development Programme (UNDP) 2000 report again result in a political catharsis and perhaps induce violent reactions? Probably not in Russia, where a more rational approach to Chernobyl prevails. But the political classes of Belarus and Ukraine

* On July 23, 2010, Belarus, Russian, and Polish news agencies, including some radio stations and TV channels, announced that this last recommendation was fulfilled by the Belarus government, which decided to repopulate 2,000 villages in the “contaminated areas.” Assuming 100 residents for one village, this would amount to about 200,000 people. It seems that preparations for this move started in about 2004, and already several thousands have come back to their old settlements. The Belarus government deserves commendation for its courage to stand up to the Chernobyl hysteria, which for years has been cultivated by Greenpeace and other Greens. Its decision brings us back to normalcy. See “Belarus Repopulating Exclusion Zone,” this issue.

have for years demonstrated a much more emotional approach. When the UNSCEAR 2000a report, documenting the low incidence of serious health hazards resulting from the Chernobyl accident, was presented to the U.N. General Assembly, the Belarus and Ukraine delegations lodged a fulminating protest. This set the stage for the Chernobyl Forum in 2002, and helped to focus its agenda.

Today, the Chernobyl rumble and emotions are beginning to settle down. In the centuries to come, the catastrophe will be remembered as a proof that nuclear power is a safe means of energy production. It even might change the thinking of the International Commission on Radiological Protection.

Zbigniew Jaworowski is a multidisciplinary scientist who has published more than 300 scientific papers, four books, and scores of popular science articles, including many in 21st Century. He has been a member of the United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR) since 1973, and served as its chairman from 1980-1982.

This article is adapted from the author's "Chernobyl Disaster and LNT," in *Dose-Response*, Vol. 8, No. 2, 2010.

References

- ACS, 2009. "Tumor markers." American Cancer Society, http://www.cancer.org/docroot/PED/content/PED_2_3X_Tumor_Markers.asp.
- L.A. Akslen and G.N. Naumov, 2008. "Tumor dormancy—from basic mechanisms to clinical practice." *Acta Pathologica, Microbiologica et Immunologica Scandinavica*, Special Issue: "Tumor Dormancy," Vol. 116, pp. 545-547.
- D. Altius, 2008. "Natural Disaster." <http://www.altiusdirectory.com/Science/natural-disaster.html>.
- H.M. ApSimon, A.J.H. Goddard, J. Wrigley, and S. Crompton, 1985. "Long-range atmospheric dispersion of radioisotopes: II. Application of the MESOS model." *Atmospheric Environment* Vol. 19, pp. 113-125.
- H.M. ApSimon and J.J.N. Wilson, 1987. "Modelling Atmospheric dispersal of the Chernobyl release across Europe." *Boundary-Layer Meteorology*, Vol. 41, pp. 123-133.
- K. Becker, 1996. "Some economical, social, and political consequences in Western Europe." Paper No. IAEA-CN-63/196. International Conference One Decade after Chernobyl: Summing up the Consequences of the Accident.
- M.L. Bell and D.L. Davis, 2001. "Reassessment of the lethal London fog of 1952: Novel indicators of acute and chronic consequences of acute exposure to air pollution." *Environmental Health Perspectives Supplements* <http://www.ehponline.org/members/2001/suppl-3/389-394bell/bell-full.html>.
- W. Benenson, J.W. Harris, H. Stocker, and H. Lutz, 2006. *Handbook of Physics*. Springer.
- A. Berrington, S.C. Darby, H.A. Weiss, and R. Doll, 2001. "100 years of observation on British radiologists: Mortality from cancer and other causes 1897-1997." *The British Journal of Radiology*, Vol. 74, pp. 507-519.
- E. Cardis et al., 2007. "The 15-country collaborative study of cancer risk among radiation workers in the nuclear industry: Estimates of radiation-related cancer risks." *Radiation Research*, Vol. 167, pp. 396-416.
- E. Cardis et al., 2006. "Estimates of the cancer burden in Europe from radioactive fallout from the Chernobyl accident." *International Journal of Cancer*, Vol. 119, pp. 1224-1235.
- E. Cardis et al., 2005. "Risk of thyroid cancer after exposure to ¹³¹I in childhood." *J of National Cancer Institute*, Vol. 97, pp. 724-732.
- W.L. Chen, Y.C. Luan, et al., 2004. "Is chronic radiation an effective prophylaxis against cancer?" *Journal of American Physicians and Surgeons*, Vol. 9, pp. 6-10.
- G.C. Christensen, 1989. "The impact of the Chernobyl accident on Norway." In *7th IRPA International Congress*, Vol. 2, pp. 1483-1486. Available at http://www2000.irpa.net/irpa7/cdrom/VOL.3/S3_106.PDF.
- R. Clarke, 1999. "Control of low-level radiation exposure: Time for a change?" *Journal of Radiological Protection*, Vol. 19, pp. 107-115.
- B.L. Cohen, 2000. "The recent cancer risk from low level radiation: A review of recent evidence." *Medical Sentinel*, Vol. 5, pp. 128-131, available at: <http://www.haciendapub.com/article50.html>.
- J.R. Davis, 2009. "Postwar relations: the long climbing from Yalta and Potsdam to Gdansk and the round table." *Polish Review*, 77 pp.
- J.R. Davis, G. Domber, M. Jarzab, P. Sowinski, et al., 2006. "Toward the Victory of Solidarity: Correspondence between the American Embassy in Warsaw and the State Department, January-September 1989" (in Polish). Instytut Studiów Politycznych PAN.
- V.R. Dhara and R. Dhara, 2002. "The Union Carbide Disaster in Bhopal: A review of Health Effects." *Archives of Environmental Health*, Vol. 57, pp. 391-404.
- J.E. Dibb, P.A. Mayewski, C.S. Buck, and S.M. Drumey, 1990. "Beta radiation from snow." *Nature*, Vol. 345, p. 25.
- P.W. Dickman, L.E. Holm, G. Lundell, and P. Hall, 2003. "Thyroid cancer risk after thyroid examination with ¹³¹I: A population-based cohort study in Sweden." *International Journal of Cancer*, Vol. 106, pp. 580-587.
- EUR, 1996. "Preliminary version of the total Caesium-137 deposition map taken from the 'Atlas of Caesium deposition on Europe after the Chernobyl accident.'" European Commission Office of Publication, Luxembourg, EUR report 16733.
- K.W. Fornalski, 2009. "What is the radiation level now in Chernobyl?" (in Polish). Press Conference of Society of Ecologists for Nuclear Energy (SEREN), Polish Press Agency, Warsaw, April 24, 2009.
- K.W. Fornalski and L. Dobrzynski, 2009. "Healthy worker effect and nuclear industry workers." *Dose-Response*, Vol. 8, No. 2: pp. 125-147.
- Forum, 2005. "Chernobyl's legacy: Health, Environmental and Socio-Economic Impacts and Recommendations to the Governments of Belarus, the Russian Federation and Ukraine," 57 pp. The Chernobyl Forum.
- Forum, 2006. "Health Effects of the Chernobyl Accident and Special Health Care Programmes." Report of the U.N. Chernobyl Forum Expert Group "Health." World Health Organization.
- N.A. Frigerio, K.F. Eckerman, and R.S. Stowe, 1973. "The Argonne Radiological Impact Program (ARIP)." Part I. "Carcinogenic Hazard from Low-level, Low-rate Radiation," 35 pp. Argonne National Laboratory.
- N.A. Frigerio and R.S. Stowe, 1976. "Carcinogenic and genetic hazard from background radiation." In *Biological and Environmental Effects of Low-Level Radiation*, Vol. 2, pp. 385-393. IAEA-SM-202/805. International Atomic Energy Agency, Vienna, Austria.
- A.W. Furmanchuk, N. Roussak, and C. Ruchti, 1993. "Occult thyroid carcinomas in the region of Minsk, Belarus. An autopsy Study of 215 patients." *Histopathology*, Vol. 23, pp. 319-325.
- M. Goldman, R.J. Catlin, and L. Anspaugh, 1987. "Health and Environmental Consequences of the Chernobyl Nuclear Power Plant Accident, 289 pp. U.S. Department of Energy.
- M. Gorbachev, 2006. "Turning point at Chernobyl." <http://www.project-syndicate.org/commentary/gorbachev3/English>.
- Greenpeace, 2006. "Chernobyl death toll grossly underestimated." April 18, 2006. Greenpeace International. <http://www.commondreams.org/cgi-bin/print.cgi?file=headlines06/0325-05.htm>.
- GUS, 1991. *Statistical Yearbook of the Republic of Poland* (in Polish). Główny Urząd Statystyczny, Warsaw, Poland.
- P. Hall, A. Mattsson, and J.D. Boice, Jr., 1996. "Thyroid cancer after diagnostic administration of iodine-131." *Radiation Research*, Vol. 145, pp. 86-92.
- H.R. Harach, K.O. Franssila, and V.M. Wasenius, 1985. "Occult papillary carcinoma of the thyroid—A 'normal' finding in Finland. A systematic study." *Cancer*, Vol. 56, pp. 531-538.
- T. Henriksen and G. Saxebol, 1988. "Fallout and radiation doses in Norway after the Chernobyl accident." *Environment International*, Special Issue: Chernobyl Accident: Regional and Global Impacts, Guest Editor, Zbigniew Jaworowski Vol. 14, pp. 157-163.
- S. Hirschberg, G. Spikerman, and R. Dones, 1998. "Severe accidents in the energy sector." Paul Scherrer Institute, Switzerland, Report No. PSI-98-16.
- L.E. Holm, P. Hall, K. Wiklund, G. Lundell, G. Berg, G. Bjelkwegren, E. Cederquist, U.B. Ericsson, L.G. Larsson, M. Lidberg, S. Lindberg, J. Tennvall, H. Wicklund, and J.J.D. Boice, 1991. "Cancer risk after iodine-131 therapy for hyperthyroidism." *Journal of the National Cancer Institute*, Vol. 83, pp. 1072-1077.
- L.E. Holm, K. Wiklund, G. Lundell, A. Bergman, G. Bjelkwegren, E. Cederquist, U.B. Ericsson, L.G. Larsson, M. Lidberg, S. Lindberg, H. Wicklund, and J.J. D. Boice, 1988. "Thyroid cancer after diagnostic doses of iodine-131: A retrospective cohort study." *Journal of the National Cancer Institute*, Vol. 80, pp. 1133-1138.
- S-Lea Hwang, 2008. "Estimates of relative risks for cancers in a population after prolonged low-dose-rate radiation exposure: A follow-up assessment from 1983 to 2005." *Radiation Research*, Vol. 170, pp. 143-148.
- IAEA, 1987. "Conclusions and Recommendations." Advisory Group Meeting on Introducing the Basic Principles of Assessment and Treatment of Radiation Injuries into the Basic and Post-Graduate Training of Medical and Paramedical Personnel.
- IAEA, 2008. "Climate Change and Nuclear Power 2008." Brochure, pp. 60. International Atomic Energy Agency <http://www.iaea.org/OurWork/ST/NE/Pess/>

- ICRP, 1959. *Recommendations of the International Commission on Radiological Protection*. Pergamon Press.
- ICRP, 1984. *Protection of the public in the event of major radiation accidents: Principles for planning*. Pergamon Press.
- ICRP, 1991. *1990 Recommendations of the International Commission on Radiological Protection*. ICRP Publication 60. Pergamon Press.
- B. Idas and J. Myhre, 1994. "Countermeasures in Norway are exaggerated" (in Norwegian). *Aftenposten*, Aug. 10, 1994.
- L.A. Ilyin, 1996. Personal communication to Z. Jaworowski, Warsaw, December 23, 1996.
- _____, 1995. *Chernobyl: Myth and Reality*. Megapolis.
- P.D. Inskip, 2001. "Thyroid cancer after radiotherapy for childhood cancer." *Medical and Pediatric Oncology*, Vol. 36, pp. 568-572.
- V.K. Ivanov, A.F. Tsyb, S. Ivanov, and V. Pokrovsky, 2004. *Medical Radiological Consequences of the Chernobyl Catastrophe in Russia*. NAUKA.
- Z. Jaworowski, 1988. "Chernobyl Proportions," Editorial. *Environ International*, Special Issue: Chernobyl Accident: Regional and Global Impacts, guest ed. Zbigniew Jaworowski, Vol. 14, pp. 69-73.
- _____, 1999. "Radiation risk and ethics." *Physics Today*, Vol. 52, pp. 24-29.
- _____, 2000. "Beneficial Radiation and Regulations." IOCON 8 8th International Conference on Nuclear Engineering, April 2-6, 2000, Baltimore.
- _____, 2002. "Ionizing radiation in the 20th Century and beyond." *Atomwirtschaft—Atomtechnik*, Vol. 47, pp. 22-27.
- _____, 2009. "Radiation Hormesis: A Remedy for Fear." *BELLE Newsletter*, Vol. 15, pp. 14-20.
- Z. Jaworowski and L. Kownacka, 1994. "Nuclear weapon and Chernobyl debris in the troposphere and lower stratosphere." *The Science of the Total Environment*, Vol. 144, pp. 201-215.
- L.E. Ketchum, 1987. "Lessons of Chernobyl: SNM members try to decontaminate the world threatened by fallout—Experts face challenge of educating public about risk and radiation." *Journal of Nuclear Medicine*, Vol. 28, pp. 933-942.
- S. Kikuchi, N. Perrier, P. Ituarte, A.E. Siperstein, Q-Y Dug, and O.H. Clark, 2004. "Latency period of thyroid neoplasia after radiation exposure." *Annals of Surgery*, Vol. 239, pp. 536-543. Available at <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1356259&rendertype=table&id=1356251-1356214>
- P. Krajewski, 1991. "Estimate of thyroid committed dose equivalents in Polish population due to iodine-131 intake after the Chernobyl catastrophe. Determination of effectiveness of thyroid blocking with sodium iodide." (in Polish). *Polish Journal of Endocrinology*, Vol. 42, pp. 189-202.
- T.D. Luckey, 2003. "Radiation hormesis overview." *RSO Magazine*, Vol. 8, pp. 22-41.
- P. McCully, 1998. "When things fall apart: The technical failures of large dams" (Chapter 4). In *Silenced Rivers: The ecology and Politics of Large dams*, p 200. South Asia Books
- M. Moosa and E.L. Mazzaferri, 1997. "Occult thyroid carcinoma." *The Cancer Journal*, Vol. 10, pp. 180-188.
- S.M.J. Mortazawi, M. Ghiassi-Neyad, P.A. Karam, T. Ikushima, A. Niroomand-rad, and J.R. Cameron, 2006. "Cancer incidence in areas with elevated levels of natural radiation." *International Journal of Low Radiation*, Vol. 2, pp. 20-27.
- K.L. Mossman, M. Goldman, F. Masse, W.A. Mills, K.J. Schiager, and R.L. Vetter, 1996. "Radiation risk in perspective." Health Physics Society Position Statement, March 1996, Vol. 12 March 1996, pp. 1-2. <http://www.physics.isu.edu/radinf/hprisk.htm>.
- G. Murphy, 2009. "A final warning for humanity—or James Lovelock?" A review of James Lovelock, *The Vanishing Face of Gaia: A Final Warning*. Basic Books, 2009, in *21st Century Science & Technology*, Vol. 22, pp. 63-64.
- J. Nauman, 1989. "Potassium iodide prophylaxis in Poland: Review of far field experience." In E. Rubery and E. Smales (eds.) *Iodine Prophylaxis Following Nuclear Accidents*, pp. 135-140. Pergamon Press.
- E.M. Parshkov, V.A. Sokolov, A.F. Tsyb, A.D. Proshin, and J.G. Barnes, 2004. "Radiation-induced thyroid cancer: What we know and what we really understand." *Int J Low Radiation*, Vol. 1, pp. 267-278.
- J.C. Philippot, 1990. "Fallout in snow." *Nature*, Vol. 348, pp. 21.
- E. Ron, 2007. "Thyroid cancer incidence among people living in areas contaminated by radiation from the Chernobyl accident." *Health Physics*, Vol. 93, pp. 502-511.
- E. Ron, J. Lubin, and A.B. Schneider, 1992. "Thyroid cancer incidence." *Nature*, Vol. 360, p. 113.
- A. Salo and J. Daglish, 1988. "Response to an accident in theory and in practice." *Environment International*, Special Issue on "Chernobyl Accident: Regional and Global Impacts," Guest Editor Zbigniew Jaworowski, Vol. 14, pp. 185-200.
- B.E. Scott, 2006. "Correspondence Re: Risk of thyroid cancer after exposure to ¹³¹I in childhood." *Journal of the National Cancer Institute*, Vol. 98, pp. 561.
- A. Spinelli and J.F. Osborne, 1991. "The effects of the Chernobyl explosion on induced abortion in Italy." *Biomedicine & Pharmacotherapy*, Vol. 45, pp. 243-247.
- G.H. Tan and H. Gharib, 1997. "Thyroid incidentalomas: Management approaches to nonpalpable nodules discovered incidentally on thyroid imaging." *Annals of Internal Medicine*, Vol. 126, pp. 226-231.
- L.S. Taylor, 1980. "Some non-scientific influences on radiation protection standards and practice." 5th International Congress of the International Radiation Protection Association.
- D. Topliss, 2004. "Thyroid incidentaloma: The ignorant in pursuit of impalpable." *Clinical Endocrinology*, Vol. 60, pp. 18-20.
- D. Trichopoulos, X. Zavitsanos, C. Koutis, P. Drogari, C. Proukakis, and E. Petridou, 1987. "The victims of Chernobyl in Greece: Induced abortions after the accident." *British Medical Journal*, Vol. 295, p. 1100.
- M. Tubiana, 1998. "The report of the French Academy of Science: 'Problems associated with the effects of low doses of ionizing radiation.'" *Journal of Radiological Protection*, Vol. 18, pp. 243-248.
- UNDP, 2002. *The Human Consequences of the Chernobyl Nuclear Accident: A strategy for Recovery*, 75 pp. United Nations Development Programme (UNDP) and the UN Children's Fund (UNICEF) with the support of the UN Office for Co-ordination of Humanitarian Affairs (OCHA) and WHO.
- UNSCEAR, 1958. *Report of the United Nations Scientific Committee on the Effects of Atomic Radiation*, 228 pp. United Nations.
- UNSCEAR, 1982. *Ionizing Radiation: Sources and Biological Effects*, 773 pp. United Nations Scientific Committee on the Effects of Atomic Radiation. United Nations.
- UNSCEAR, 1988. *Sources, Effects and Risks of Ionizing Radiation. Report of the United Nations Scientific Committee on the Effects of Atomic Radiation*, 647 pp. United Nations.
- UNSCEAR, 1993. *Sources and Effects of Ionizing Radiation*, 922 pp. United Nations Scientific Committee on the Effects of Atomic Radiation. United Nations.
- UNSCEAR, 1994. "Annex B: Adaptive responses to radiation in cells and organisms." In *Sources and Effects of Ionizing Radiation. Report of the United Nations Scientific Committee on the Effects of Atomic Radiation*, pp 185-272. United Nations.
- UNSCEAR, 1998. *Exposures from man-made radiation*. Report of United Nations Scientific Committee on the Effects of Atomic Radiation, 130 pp. United Nations.
- UNSCEAR, 2000a. *Sources and Effects of Ionizing Radiation. United Nations Scientific Committee on the Effects of Atomic Radiation*, UNSCEAR 2000, Report to the General Assembly. United Nations.
- UNSCEAR, 2000b. *Sources and Effects of Ionizing Radiation*. United Nations Scientific Committee on the Effects of Atomic Radiation UNSCEAR 2000, Report to the General Assembly. "Annex J: Exposures and Effects of the Chernobyl Accident," pp. 451-566. United Nations.
- UNSCEAR, 2001. *Hereditary Effects of Radiation*. "Scientific annex" of UNSCEAR 2001 report to the General Assembly, 224 pp. United Nations Scientific Committee on the Effects of Atomic Radiation. United Nations.
- UNSCEAR, 2008. *Health effects due to radiation from the Chernobyl accident*. Draft report A/AC.82/R.673, 220 pp. United Nations Scientific Committee on the Effects of Atomic Radiation. United Nations.
- J. Vidal, 2006. "UN accused of ignoring 500,000 Chernobyl deaths." *The Guardian* (U.K.) March 25, 2006. <http://www.commondreams.org/cgi-bin/print.cgi?file=/headlines06/0325-05.htm>.
- G. Walinder, 1995. "Has radiation protection become a health hazard?" The Swedish Nuclear Training & Safety Center.
- E.W. Webster, 1993. "Hormesis and radiation protection." *Investigative Radiology*, Vol. 28, pp. 451-453.
- L. Wei, Y. Zha, Z. Tao, W. He, D. Chen, and Y. Yuan, 1990. "Epidemiological investigation of radiological effects in high background radiation areas of Yangjiang, China." *Journal of Radiation Research*, Vol. 31, pp. 119-136.
- R.A. Weinberg, 2008. "The many faces of tumor dormancy." *Acta pathologica, Microbiologica, et Immunologica Scandinavica*. Special Issue: Tumor Dormancy, Vol. 116, pp. 548-551. <http://www3.interscience.wiley.com/cgi-bin/fulltext/121415236/PDFSTART>.
- WNA, 2009. *The Hazards of Using Energy: Some energy-related accidents since 1977*. World Nuclear Association. <http://www.world-nuclear.org/info/inf06app.htm>.
- S. Yi, 1998. "The World's Most Catastrophic Dam Failures. The August 1975 collapse of the Baqiao and Shimantan dams." In: D. Qing, J. Thiboleau, and P.B. Williams (eds.) *The River Dragon Has come! The Three Gorges Dam and the Fate of China's Yangtze River and its People*, 240 pp.